



**36% BEEF GROWER SUPPLEMENT PELLET**

Feed to beef cattle

**MEDICATED**

Cattle being fed in confinement for slaughter; For the prevention and control of coccidiosis due to Eimeria bovis and Eimeria zuernii

**ACTIVE DRUG INGREDIENT**

Monensin (as Monensin sodium)..... 95 g/ton

This feed contains added selenium at 1.55 mg/kg.

**GUARANTEED ANALYSIS**

CRUDE PROTEIN*	(Min)	35.4 %	SALT	(Min)	2.10 %
CRUDE FAT	(Min)	2.6 %	SALT	(Max)	2.50 %
CRUDE FIBER	(Max)	6.3 %	SODIUM	(Min)	0.68 %
CALCIUM	(Min)	3.55 %	SODIUM	(Max)	1.08 %
CALCIUM	(Max)	3.95 %	POTASSIUM	(Min)	0.89 %
PHOSPHORUS	(Min)	0.60 %	VITAMIN A	(Min)	11,111 IU/LB

\* (Includes not more than 12.50 % equivalent crude protein from non-protein nitrogen).

**INGREDIENTS**

Processed Grain By- Products, Plant Protein Products, Calcium Carbonate, Urea, Sodium Chloride, Potassium Chloride, Magnesium Oxide, AMERIBOND PELLET BINDER, Calcium Carbonate, Sodium Selenite, Mineral Oil, Zinc Hydroxychloride, Vitamin E Supplement, Zinc Proteinat, Manganese Hydroxychloride, Copper Sulfate, Manganese Proteinat, Zinc Sulfate, Ethylenediamine Dihydroiodide, Manganese Sulfate, Soybean Oil, Vitamin A Supplement, Basic Copper Chloride, Vitamin D Supplement, Copper Proteinat, Cobalt Carbonate.

**DIRECTIONS FOR USE:**

Feed this Shur-Gain Beef supplement at 4.26 lb/hd/day along with forages and grains to cattle being fed for slaughter.

**CAUTION:** A withdrawal time has not been established for preruminating calves. Do not use in calves to be processed for veal. Do not allow horses or other equines access to formulation containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin medicated cattle and goat feed is safe for use in cattle and goats only. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of Monensin recommended in the feeding directions as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and amount of refusals fed should be taken into consideration to prevent Monensin overdosing.

**LIMITATIONS:** For prevention and control of coccidiosis, feed at a rate of 0.14 to 0.42 milligram per pound of body weight per day, depending on severity of challenge, up to 200 milligrams per head per day. During first 5 days of feeding, cattle should receive no more than 100 milligrams per day in not less than 1 pound of feed.

**THIS FEED CONTAINS UREA!** Use with care particularly when feeding with silage containing added urea or other sources of non-protein nitrogen.

1. Directions for use must be carefully followed. 2. Do not use this feed in association with another grain ration, supplement, or premix containing supplemental selenium.

Manufactured by: Nutreco USA, Inc.  
P.O. Box 69, 3422 Dutch Hollow Rd.,  
Strykersville, New York 14145

50 lb

**10870735**

**REMINDER:** Because individual results from the use of this product may vary due to management, environment, genetic, health and sanitation differences, Nutreco USA, Inc. and its distribution system do not warrant or guarantee individual results.

**MEDICATED**