



HI DEF P/C SHOW BEEF SUPPLEMENT TEXTURED

MEDICATED

Cattle being fed in confinement for slaughter; For the prevention and control of coccidiosis due to Eimeria bovis and Eimeria zuernii

ACTIVE DRUG INGREDIENT

Monensin (as Monensin sodium)..... 17 g/ton

This feed contains added selenium at 0.26 mg/kg.

GUARANTEED ANALYSIS

CRUDE PROTEIN	(Min)	13.2 %	SALT	(Min)	0.25 %
CRUDE FAT	(Min)	3.8 %	SALT	(Max)	0.65 %
CRUDE FIBER	(Max)	11.7 %	SODIUM	(Min)	0.05 %
CALCIUM	(Min)	0.42 %	SODIUM	(Max)	0.45 %
CALCIUM	(Max)	0.82 %	POTASSIUM	(Min)	0.35 %
PHOSPHORUS	(Min)	0.11 %	VITAMIN A	(Min)	2,762 IU/LB

INGREDIENTS

Grain Products, Roughage Products, Plant Protein Products, Animal Fat, Molasses Products, Calcium Carbonate, Sodium Selenite, Mineral Oil.

DIRECTIONS FOR USE:

Feed this custom beef feed according to the feeding program provided by the Trouw Nutrition sales representative or farm Nutritional Consultant.

Each pound of this feed contains 08.7 mg Monensin. Feed continuously to beef cattle at a rate of 5.8 lbs/head/day to 54.7 lbs/head/day to provide not less than 50 mg/head/day nor more than 480 mg/head/day Monensin.

CAUTION: A withdrawal time has not been established for prerinuating calves. Do not use in calves to be processed for veal. Do not allow horses or other equines access to formulation containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin medicated cattle and goat feed is safe for use in cattle and goats only. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of Monensin recommended in the feeding directions as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and amount of refusals fed should be taken into consideration to prevent Monensin overdosing.

LIMITATIONS: For prevention and control of coccidiosis, feed at a rate of 0.14 to 0.42 milligram per pound of body weight per day, depending on severity of challenge, up to 200 milligrams per head per day. During first 5 days of feeding, cattle should receive no more than 100 milligrams per day in not less than 1 pound of feed.

1. Directions for use must be carefully followed. 2. Do not use this feed in association with another grain ration, supplement, or premix containing supplemental selenium.

Manufactured by: Nutreco USA, Inc.
P.O. Box 69, 3422 Dutch Hollow Rd.,
Strykersville, New York 14145

50 lb

13314675

REMINDER: Because individual results from the use of this product may vary due to management, environment, genetic, health and sanitation differences, Nutreco USA, Inc. and its distribution system do not warrant or guarantee individual results.

MEDICATED