



HI DEF 18% SHOW SHEEP FEED TEXTURED

MEDICATED

For the prevention of coccidiosis caused by Eimeria ovina, Eimeria crandalis, Eimeria ovinoidalis (Eimeria ninakohlyakimovae), Eimeria parva and Eimeria intricata in sheep maintained in confinement.

ACTIVE DRUG INGREDIENT

Lasalocid 27 g/ton

This feed contains added selenium at 0.35 mg/kg.

GUARANTEED ANALYSIS

CRUDE PROTEIN*	(Min)	17.8 %	SALT	(Min)	0.85 %
CRUDE FAT	(Min)	3.7 %	SALT	(Max)	1.25 %
CRUDE FIBER	(Max)	5.0 %	SODIUM	(Min)	0.46 %
CALCIUM	(Min)	1.23 %	SODIUM	(Max)	0.86 %
CALCIUM	(Max)	1.63 %	SELENIUM	(Min)	0.35 ppm
PHOSPHORUS	(Min)	0.40 %	VITAMIN A	(Min)	5,345 IU/LB
* (Includes not more than 0.99 % equivalent crude protein from non-protein nitrogen).					

INGREDIENTS

Grain Products, Plant Protein Products, Molasses Products, Calcium Carbonate, Mono Dicalcium Phosphate, Calcium Carbonate, Salt, Magnesium Oxide, Magnesium Potassium Sulfate, Potassium Chloride, Zinc Sulfate, Processed Grain By-Products, Manganese Sulfate, Mineral Oil, Sodium Selenite, Vitamin E Supplement, Vitamin A Supplement, Calcium Iodate, Vitamin D Supplement, Cobalt Carbonate, Sodium Bicarbonate, Ammonium Chloride, Sodium Chloride, Calcium Carbonate, Vitamin E Supplement, Vitamin A supplement, Vitamin D Supplement, Mineral Oil.

DIRECTIONS FOR USE:

Feed this custom sheep feed according to the feeding program provided by the Trouw Nutrition sales representative or farm Nutritional Consultant.

Each pound of this feed contains 13.6 mg Lasalocid. Feed continuously to sheep at a rate of 1.1 lbs/head/day to 5.1 lbs/head/day to provide not less than 15 mg/head/day nor more than 75 mg/head/day Lasalocid depending on bodyweight.

CAUTION: The safety of Lasalocid for use in unapproved species has not been established. Do NOT allow horses or other equines access to feeds containing Lasalocid as ingestion may be fatal. For Type A and Type B feeds (cattle and sheep): mix thoroughly with grain and/or roughage products prior to feeding. Feeding undiluted or having mixing errors may result in excess Lasalocid concentration which could be fatal to cattle and sheep. In addition, inadequate mixing (recirculation or agitation) of liquid Type B feed may result in an excess Lasalocid concentration which could be fatal to cattle and sheep.

LIMITATIONS: In Type C feeds; for sheep maintained in confinement; feed continuously in complete feed to provide not less than 15 mg nor more than 70 mg of lasalocid sodium activity per head per day depending on body weight.

1. Directions for use must be carefully followed. 2. Do not use this feed in association with another grain ration, supplement, or premix containing supplemental selenium.

Manufactured by: Nutreco USA, Inc.
P.O. Box 69, 3422 Dutch Hollow Rd.,
Strykersville, New York 14145

50 lb

12679935

REMINDER: Because individual results from the use of this product may vary due to management, environment, genetic, health and sanitation differences, Nutreco USA, Inc. and its distribution system do not warrant or guarantee individual results.

MEDICATED