

AGRI-BEST 24/30 LAMB MILK REPLACER MEDICATED

For the prevention of coccidiosis in young sheep caused by *Eimeria ovinoidalis*, *E. crandallis*, *E. parva*, and *E. bakuensis*.

Thoroughly mix Decco into the ration at a rate to provide decoquinatate at a daily dose of 22.7 mg/100 lbs (0.5 mg/kg) of body weight. Feed for at least 28 days during periods of coccidiosis exposure or when experience indicates that coccidiosis is likely to be a hazard.

ACTIVE DRUG INGREDIENT

Decoquinatate.....18.16 grams/ton

GUARANTEED ANALYSIS

Crude Protein, not less than.....	24.00%
Crude Fat, not less.....	30.00%
Crude Fiber, not more than.....	0.15%
Calcium, min.....	0.75%
Calcium, max.....	1.25%
Phosphorus, min.....	0.70%
Copper, min2 ppm
Copper, max	6 ppm
Selenium, min	0.3 ppm
Lactose, max	25.00%
Vitamin A, not less than.....	30,000 IU/lb
Vitamin D3, not less than.....	10,000 IU/lb
Vitamin E, not less than.....	200 IU/lb
Total Lactic Acid Bacteria	2 billion CFU*/lb
.....	*CFU (Colony Forming Units)

INGREDIENTS

Dried Whey Protein Concentrate, Dried Whey, Animal and Vegetable Fat (Preserved with BHT), Dried Skimmed Milk, Dextrose, Lecithin, Dried Milk Protein, Dicalcium Phosphate, Calcium Carbonate, Magnesium Sulfate, L-Lysine, dl-Methionine, Vitamin A Acetate, Vitamin D₃ Supplement, Vitamin E Supplement, Manganese Proteinatate, Zinc Proteinatate, Iron Proteinatate, Selenium Yeast, Dried Brewer's Yeast, Dried *Lactobacillus acidophilus* Fermentation Product, Dried *Lactobacillus casei* Fermentation Product, Dried *Bifidobacterium bifidum* Fermentation Product, Dried *Streptococcus diacetylactis* Fermentation Product, Dried *Bifidobacterium licheniformis* Fermentation Product, Dried *Bifidobacterium subtilis* Fermentation Product, Vitamin B₁₂ Supplement, Thiamine Mononitrate, Ascorbic Acid, Biotin, Riboflavin, d-Calcium Pantothenate, Niacinamide, Choline Chloride, Menadione Sodium Bisulfite Complex (Source of Vitamin K₃), Folic Acid, Pyridoxine Hydrochloride, Calcium Iodate, Zinc Sulfate, Iron Sulfate, Manganese Sulfate, Copper Sulfate, Cobalt Sulfate, Sodium Silico Aluminate, Polyoxyethylene Glycol (400) Mono and Dioleates, Artificial Flavor.

Warning: Do not feed to sheep producing milk for food

**Manufactured by
STRAUSS FEEDS LLC
Watertown, WI 53098**

NET WEIGHT 25 LB. (11.33 KG.)

FEEDING AND MIXING INSTRUCTIONS

APPLICATION

Agri-Best Lamb Milk Replacer is a fully formulated complete first feed for lambs and goat kids. It provides the entire nutritional requirements of the baby lamb or kid from its second day of life through to weaning.

FEEDING INSTRUCTIONS: LAMBS

1. Ensure that the baby lamb receives colostrum during its first 12 hours of life.
2. For best results, mix Agri-Best Lamb Milk Replacer with warm water (110° F), then cool to room temperature. Offer ad libitum to lambs through a suitable nipple feeder. Milk should be available at all times.
3. For hand feeding with a nipple bottle or restricted feeding offer warm milk at 100° F. It is recommended to offer 3 to 6 feedings per day of warm milk during the first two weeks of life and thereafter 2 feedings per day. Never offer more than 1/2 pint per feeding.
4. Provide good quality lamb creep at 3 to 6 days of age and restrict hay intake to the amount a lamb will eat in an hour.
5. Wean lambs at a minimum of 30 days and weighing at least 25 pounds.

MIXING INSTRUCTIONS

Mix at a rate of 2 lbs. per gallon of warm water (110° F) or one (1) 4 ounce cup of Agri-Best Lamb Milk Replacer with two (2) cups of warm water. Milk will stay fresh up to 3 days due to acidification.

FEEDING INSTRUCTION: GOAT KIDS

1. Restricted feeding: For the first few days, feed a limited quantity, up to 1 pint per feeding up to 3 to 4 times a day. As kids get older reduce the number of feedings per day. Best results are achieved feeding milk warm (98° F).
2. Ad libitum feeding: This method works well for raising larger groups of kids that have been removed from the mother soon after birth. It is the recommended systems when a number of kids are being taken away, as with a milking herd.

MIXING INSTRUCTIONS

Mix at a rate of 1.5 lb. per gallon of water (3 ounces per pint).

1. Warm Restricted Feeding: Mix powder in 1/2 quantity of water at 110° F and then dilute to volume and feed at 98° F.
2. For Cold Ad Lib Feeding: Mix powder in 1/3 quantity of water at 110° F and then dilute to volume and feed at room temperature.

*One level measuring cup equals 4 ounces of dry Agri-Best Lamb Milk Replacer.