



OPTIVIA 22% CALF STARTER TEXTURED

MEDICATED

For the prevention and control of coccidiosis due to Eimeria bovis and Eimeria zuernii in calves (excluding veal calves).

ACTIVE DRUG INGREDIENT

Monensin (as Monensin sodium)..... 45 g/ton

This feed contains added selenium at 0.30 mg/kg.

GUARANTEED ANALYSIS

CRUDE PROTEIN	(Min)	21.7 %	CALCIUM	(Max)	1.34 %
CRUDE FAT	(Min)	4.6 %	PHOSPHORUS	(Min)	0.51 %
CRUDE FIBER	(Max)	6.1 %	SELENIUM	(Min)	0.30 ppm
CALCIUM	(Min)	0.94 %	VITAMIN A	(Min)	7,217 IU/LB

INGREDIENTS

Grain Products, Molasses Products, Processed Grain ByProducts, Soybean Oil, Roughage Products, Whey Powder, Plant Protein Products, Hydrolyzed Animal and Vegetable Fat (Preserved with BHT), Magnesium Oxide, Calcium Propionate, Urea Formaldehyde Resin, Kaolin, Calcium Sulphate, Silicon Dioxide, Natural and Artificial Flavors.

DIRECTIONS FOR USE:

Feed this custom dairy heifer grain according to the feeding program provided by the Shur-Gain sales representative or farm Nutritional Consultant, or at a rate of 2.20 lbs. to 8.82 lbs. per head per day to provide not less than 50 nor more than 200 mg monensin per head per day depending on the body weight of the animals and not less than 1 lb of feed. Feed continuously during periods of exposure to coccidia or when coccidiosis is likely to be hazard.

CAUTION: Do not allow horses or other equines access to formulation containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin medicated cattle and goat feed is safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Do not feed undiluted. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Do not exceed the levels of Monensin recommended by the feeding directions, as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and amount of refusals fed should be taken into consideration to prevent Monensin overdosing.

WARNING: A withdrawal time has not been established for prerininating calves. Do not use in calves to be processed for veal.

LIMITATIONS: For calves, excluding veal calves, feed at a rate of 0.14 to 1.0 milligrams Monensin per pound of body weight per day, depending upon the severity of challenge, up to maximum of 200 milligrams per head per day.

Manufactured by: Nutreco USA, Inc.
P.O. Box 69, 3422 Dutch Hollow Rd.,
Strykersville, New York 14145

Net Weight on Bag or Bulk Invoice

10873885

REMINDER: Because individual results from the use of this product may vary due to management, environment, genetic, health and sanitation differences, Nutreco USA, Inc. and its distribution system do not warrant or guarantee individual results.

MEDICATED