



TROUW 16% HEIFER GROWER RATION MASH

MEDICATED

Mature beef cows in confinement: For improved feed efficiency. For prevention and control of coccidiosis caused by Eimeria bovis and Eimeria zuernii.

ACTIVE DRUG INGREDIENT

Monensin Sodium 80 g/ton

This feed contains added selenium at 0.60 mg/kg.

GUARANTEED ANALYSIS

CRUDE PROTEIN	(Min)	16.0 %	CALCIUM	(Max)	1.25 %
CRUDE FAT	(Min)	4.3 %	PHOSPHORUS	(Min)	0.67 %
CRUDE FIBER	(Max)	6.6 %	SELENIUM	(Min)	0.60 ppm
CALCIUM	(Min)	0.85 %	VITAMIN A	(Min)	3,333 IU/LB

INGREDIENTS

Processed Grain By- Products, Grain Products, Molasses Products, Calcium Carbonate, Sodium Chloride, Calcium Carbonate, Sodium Selenite, Mineral Oil, Zinc Hydroxychloride, Vitamin E Supplement, Zinc Proteinate, Manganese Hydroxychloride, Copper Sulfate, Manganese Proteinate, Zinc Sulfate, Ethylenediamine Dihydroiodide, Manganese Sulfate, Soybean Oil, Vitamin A Supplement, Basic Copper Chloride, Vitamin D Supplement, Copper Proteinate, Cobalt Carbonate.

DIRECTIONS FOR USE:

Feed this custom dairy heifer grain according to the feeding program provided by the Shur-Gain sales representative or farm Nutritional Consultant, or at a rate of 1.25 lbs. to 4.99 lbs. per head per day to provide not less than 50 nor more than 200 mg monensin per head per day depending on the body weight of the animals and not less than 1 lb of feed. Feed continuously during periods of exposure to coccidia or when coccidiosis is likely to be hazard.

CAUTION: A withdrawal time has not been established for preruminating calves. Do not use in calves to be processed for veal. Do not allow horses or other equines access to formulation containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin medicated cattle and goat feed is safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of Monensin recommended in the feeding directions as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and amount of refusals fed should be taken into consideration to prevent Monensin overdosing.

LIMITATIONS: Feed to mature reproducing beef cows. Feed as supplemental feed, either hand-fed in a minimum of 1 pound of feed or mixed in a total ration. For improved feed efficiency, feed continuously at a rate of 50 to 200 milligrams Monensin per head per day. For prevention and control of coccidiosis, feed at a rate of 0.14 to 0.42 milligram per pound of body weight per day, depending on severity of challenge, up to a maximum of 200 milligrams per head per day. During first 5 days of feeding, cattle should receive no more than 100 milligrams per head per day.

Manufactured by: Nutreco USA, Inc.
P.O. Box 69, 3422 Dutch Hollow Rd.,
Strykersville, New York 14145

Net Weight on Bag or Bulk Invoice

10870485

REMINDER: Because individual results from the use of this product may vary due to management, environment, genetic, health and sanitation differences, Nutreco USA, Inc. and its distribution system do not warrant or guarantee individual results.

MEDICATED