



TROUW 19% CALF STARTER RATION TEXTURED

MEDICATED

For the prevention and control of coccidiosis due to Eimeria bovis and Eimeria zuernii in calves (excluding veal calves).

ACTIVE DRUG INGREDIENT

Monensin (as Monensin sodium)..... 46 g/ton

This feed contains added selenium at 0.36 mg/kg.

GUARANTEED ANALYSIS

CRUDE PROTEIN*	(Min)	19.2 %	CALCIUM	(Max)	1.24 %
CRUDE FAT	(Min)	3.3 %	PHOSPHORUS	(Min)	0.53 %
CRUDE FIBER	(Max)	6.2 %	SELENIUM	(Min)	0.36 ppm
CALCIUM	(Min)	0.84 %	VITAMIN A	(Min)	5,308 IU/LB

* (Includes not more than 0.22 % equivalent crude protein from non-protein nitrogen).

INGREDIENTS

Grain Products, Processed Grain By- Products, Molasses Products, Plant Protein Products, Soybean Oil, Roughage Products, Calcium Carbonate, Sodium Chloride, Whey Powder, Calcium Propionate, Hydrolyzed Animal and Vegetable Fat (Preserved with BHT), Magnesium Oxide, AMERIBOND PELLET BINDER, Vitamine E Supplement, Selenium yeast, Calcium Carbonate, Sodium Selenite, Mineral Oil, Magnesium Oxide, Calcium Carbonate, Vitamin E Supplement, Vitamin A supplement, Vitamin D Supplement, Mineral Oil, Urea Formaldehyde Resin, Kaolin, Calcium Sulphate, Silicon Dioxide, Zinc Hydroxychloride, Vitamin E Supplement, Copper Sulfate, Zinc Proteinate, Manganese Hydroxychloride, Natural and Artificial Flavors, Manganese Proteinate, Zinc Sulfate, Ethylenediamine Dihydroiodide, Manganese Sulfate, Vitamin A Supplement, Basic Copper Chloride, Vitamin D Supplement, Copper Proteinate, Cobalt Carbonate.

DIRECTIONS FOR USE:

Feed this calf feed according to the feeding program provided by the Trouw Nutrition sales representative or farm Nutritional Consultant.

Each pound of this feed contains 23.2 mg Monensin. Feed to calves at a rate to provide not less than 0.14 mg nor more than 1.0 mg Monensin, per pound of bodyweight per day, up to 200 mg per head per day, depending on the severity of challenge.

CAUTION: Do not allow horses or other equines access to formulation containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin medicated cattle and goat feed is safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Do not feed undiluted. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Do not exceed the levels of Monensin recommended by the feeding directions, as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and amount of refusals fed should be taken into consideration to prevent Monensin overdosing.

WARNING: A withdrawal time has not been established for preruminating calves. Do not use in calves to be processed for veal.

LIMITATIONS: For calves, excluding veal calves, feed at a rate of 0.14 to 1.0 milligrams Monensin per pound of body weight per day, depending upon the severity of challenge, up to maximum of 200 milligrams per head per day.

THIS FEED CONTAINS UREA! Use with care particularly when feeding with silage containing added urea or other sources of non-protein nitrogen. 1. Directions for use must be carefully followed. 2. Do not use this feed in association with another grain ration, supplement, or premix containing supplemental selenium.

Manufactured by: Nutreco USA, Inc.
P.O. Box 69, 3422 Dutch Hollow Rd.,
Strykersville, New York 14145

50 lb

11731685

REMINDER: Because individual results from the use of this product may vary due to management, environment, genetic, health and sanitation differences, Nutreco USA, Inc. and its distribution system do not warrant or guarantee individual results.

MEDICATED