



**TROUW 22% STARTER/GROWER RATION PELLETT**

Feed to dairy and beef replacement heifers.

**MEDICATED**

For the prevention and control of coccidiosis due to Eimeria bovis and Eimeria zuernii in calves (excluding veal calves).

**ACTIVE DRUG INGREDIENT**

Monensin (as Monensin sodium)..... 55 g/ton

This feed contains added selenium at 0.35 mg/kg.

**GUARANTEED ANALYSIS**

CRUDE PROTEIN	(Min)	21.3 %	CALCIUM	(Max)	1.21 %
CRUDE FAT	(Min)	2.4 %	PHOSPHORUS	(Min)	0.76 %
CRUDE FIBER	(Max)	8.5 %	SELENIUM	(Min)	0.35 ppm
ADF	(Max)	11.5 %	VITAMIN A	(Min)	3,950 IU/LB
CALCIUM	(Min)	0.81 %			

**INGREDIENTS**

Processed Grain By- Products, Plant Protein Products, Grain Products, Roughage Products, Calcium Carbonate, XLA-20, Sodium Chloride, AMERIBOND PELLETT BINDER, Natural and Artificial Flavors, Vitamine E Supplement, Selenium yeast, Calcium Carbonate, Sodium Selenite, Mineral Oil, Calcium Carbonate, Vitamin E Supplement, Vitamin A supplement, Vitamin D Supplement, Mineral Oil, Yeast active dehydrated, Zinc Hydroxychloride, Vitamin E Supplement, Zinc Proteinat, Manganese Hydroxychloride, Manganese Proteinat, Zinc Sulfate, Ethylenediamine Dihydroiodide, Manganese Sulfate, Soybean Oil, Animal Fat, Vitamin A Supplement, Basic Copper Chloride, Vitamin D Supplement, Copper Proteinat, Cobalt Carbonate.

**DIRECTIONS FOR USE:**

Each pound of this Trouw Nutrition calf feed contains 27.45 mg monensin. Feed at a rate to provide 0.14-1.0 mg monensin per pound of bodyweight per head per day, up to 7.29 lbs. per head per day to provide no more than 200 mg monensin per head per day depending on the severity of challenge. Feed continuously during periods of exposure to coccidia or when coccidiosis is likely to be hazard.

**CAUTION:** Do not allow horses or other equines access to formulation containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin medicated cattle and goat feed is safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Do not feed undiluted. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Do not exceed the levels of Monensin recommended by the feeding directions, as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and amount of refusals fed should be taken into consideration to prevent Monensin overdosing.

**WARNING:** A withdrawal time has not been established for preruminating calves. Do not use in calves to be processed for veal.

**LIMITATIONS:** For calves, excluding veal calves, feed at a rate of 0.14 to 1.0 milligrams Monensin per pound of body weight per day, depending upon the severity of challenge, up to maximum of 200 milligrams per head per day.

Manufactured by: Nutreco USA, Inc.  
P.O. Box 69, 3422 Dutch Hollow Rd.,  
Strykersville, New York 14145

50 lb

12916815

REMINDER: Because individual results from the use of this product may vary due to management, environment, genetic, health and sanitation differences, Nutreco USA, Inc. and its distribution system do not warrant or guarantee individual results.

**MEDICATED**