



# TN Heifer DD Rumensin 4oz



# TN Heifer DD Rumensin 4oz



## SGHEFRUM MEDICATED

For the prevention and control of coccidiosis due to Eimeria bovis and Eimeria zuernii in pasture cattle (slaughter, stocker and feeder, dairy and beef replacement). For increased rate of weight gain.

### ACTIVE DRUG INGREDIENT

Monensin (as Monensin sodium). 794 g/ton  
Guaranteed Analysis

SALT	Minimum	12	%
CALCIUM	Minimum	28.36	%
MAGNESIUM	Minimum	5.09	%
POTASSIUM	Minimum	1.69	%
SALT	Minimum	12	%
SODIUM	Minimum	4.67	%
CHLORIDE	Minimum	8.78	%
ZINC	Minimum	1,730	PPM
IRON	Minimum	741	ppm
MANGANESE	Minimum	1,255	PPM
COPPER	Minimum	649	PPM
COBALT	Minimum	24	PPM
IODINE	Minimum	238	PPM
SELENIUM	Minimum	18	PPM
vitamin A	Minimum	79,450	IU/LB
vitamin D	Minimum	36,320	IU/LB
vitamin E	Minimum	987	IU/LB
ORGANIC SELE.	Minimum	10	PPM

### Ingredient List

Calcium carbonate, Salt, Magnesium Oxide, Potassium Chloride, Intellibond micro 5, Mineral oil, monensin, vitamin E supplement, Selenium Yeast, Tri Basic Copper, Manganese Oxide, Sodium Selenite, vitamin A Acetate, Calcium Iodate, Vitamin D3 supplement, Ethylene Diamine Dihydro Iodide.

**CAUTION:** A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal. Do not allow horses or other equines access to formulation containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin medicated cattle and goat feed is safe for use in cattle and goats only. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of Monensin recommended in the feeding directions as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and amount of refusals fed should be taken into consideration to prevent Monensin overdosing.

**LIMITATIONS:** For increased rate of weight gain, feed at a rate of 50 to 200 milligrams Monensin per head per day in not less than 1 pound of feed or, after the 5th day, feed at a rate of 400 milligrams per head per day every other day in not less than 2 pounds of feed. For prevention and control of coccidiosis, feed at a rate of 0.14 to 0.42 milligram per pound of body weight per day, depending on severity of challenge, up to 200 milligrams per head per day. During first 5 days of feeding, cattle should receive no more than 100 milligrams per day in not less than 1 pound of feed.

**NET WT 50 LBS (22.67) KG**

SGHEFRUM1

Manufactured for: Trouw Nutrition  
3422 Dutch Hollow Rd  
Strykersville, NY, 14145  
BATCH 37385

10/23/2020



SGHEFRUM1

**MEDICATED**

## SGHEFRUM MEDICATED

For the prevention and control of coccidiosis due to Eimeria bovis and Eimeria zuernii in pasture cattle (slaughter, stocker and feeder, dairy and beef replacement). For increased rate of weight gain.

### ACTIVE DRUG INGREDIENT

Monensin (as Monensin sodium). 794 g/ton  
Guaranteed Analysis

SALT	Minimum	12	%
CALCIUM	Minimum	28.36	%
MAGNESIUM	Minimum	5.09	%
POTASSIUM	Minimum	1.69	%
SALT	Minimum	12	%
SODIUM	Minimum	4.67	%
CHLORIDE	Minimum	8.78	%
ZINC	Minimum	1,730	PPM
IRON	Minimum	741	ppm
MANGANESE	Minimum	1,255	PPM
COPPER	Minimum	649	PPM
COBALT	Minimum	24	PPM
IODINE	Minimum	238	PPM
SELENIUM	Minimum	18	PPM
vitamin A	Minimum	79,450	IU/LB
vitamin D	Minimum	36,320	IU/LB
vitamin E	Minimum	987	IU/LB
ORGANIC SELE.	Minimum	10	PPM

### Ingredient List

Calcium carbonate, Salt, Magnesium Oxide, Potassium Chloride, Intellibond micro 5, Mineral oil, monensin, vitamin E supplement, Selenium Yeast, Tri Basic Copper, Manganese Oxide, Sodium Selenite, vitamin A Acetate, Calcium Iodate, Vitamin D3 supplement, Ethylene Diamine Dihydro Iodide.

**CAUTION:** A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal. Do not allow horses or other equines access to formulation containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin medicated cattle and goat feed is safe for use in cattle and goats only. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of Monensin recommended in the feeding directions as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and amount of refusals fed should be taken into consideration to prevent Monensin overdosing.

**LIMITATIONS:** For increased rate of weight gain, feed at a rate of 50 to 200 milligrams Monensin per head per day in not less than 1 pound of feed or, after the 5th day, feed at a rate of 400 milligrams per head per day every other day in not less than 2 pounds of feed. For prevention and control of coccidiosis, feed at a rate of 0.14 to 0.42 milligram per pound of body weight per day, depending on severity of challenge, up to 200 milligrams per head per day. During first 5 days of feeding, cattle should receive no more than 100 milligrams per day in not less than 1 pound of feed.

**NET WT 50 LBS (22.67) KG**

SGHEFRUM1

Manufactured for: Trouw Nutrition  
3422 Dutch Hollow Rd  
Strykersville, NY, 14145  
BATCH 37385

10/23/2020



SGHEFRUM1

**MEDICATED**

# TN Heifer 4oz DD

SGHEIFER

## Guaranteed Analysis

SALT	Minimum	12 %
CALCIUM	Minimum	28.53 %
MAGNESIUM	Minimum	5.09 %
POTASSIUM	Minimum	1.69 %
SALT	Minimum	12 %
SODIUM	Minimum	4.67 %
CHLORIDE	Minimum	8.78 %
ZINC	Minimum	1,730 PPM
IRON	Minimum	745 ppm
MANGANESE	Minimum	1,255 PPM
COPPER	Minimum	649 PPM
COBALT	Minimum	24 PPM
IODINE	Minimum	238 PPM
SELENIUM	Minimum	18 PPM
vitamin A	Minimum	79,450 IU/LB
vitamin D	Minimum	36,320 IU/LB
vitamin E	Minimum	987 IU/LB
ORGANIC SELE	Minimum	10 PPM

### Ingredient List

Calcium carbonate, Salt, Magnesium Oxide, Potassium Chloride, Intellibond micro 5, Mineral oil, vitamin E supplement, Selenium Yeast, Tri Basic Copper, Manganese Oxide, Sodium Selenite, vitamin A Acetate, Calcium Iodate, Vitamin D3 supplement, Ethylene Diamine Dihydro Iodide.

**NET WT 50 LBS (22.67) KG**

SGHEIFER1

**Manufactured for: Trouw Nutrition**

**3422 Dutch Hollow Rd**

**Strykersville, NY, 14145**

10/23/2020



SGHEIFER1

CONSULT YOUR FEED NUTRITIONIST FOR FEEDING RATES.

# TN Heifer 4oz DD

SGHEIFER

## Guaranteed Analysis

SALT	Minimum	12 %
CALCIUM	Minimum	28.53 %
MAGNESIUM	Minimum	5.09 %
POTASSIUM	Minimum	1.69 %
SALT	Minimum	12 %
SODIUM	Minimum	4.67 %
CHLORIDE	Minimum	8.78 %
ZINC	Minimum	1,730 PPM
IRON	Minimum	745 ppm
MANGANESE	Minimum	1,255 PPM
COPPER	Minimum	649 PPM
COBALT	Minimum	24 PPM
IODINE	Minimum	238 PPM
SELENIUM	Minimum	18 PPM
vitamin A	Minimum	79,450 IU/LB
vitamin D	Minimum	36,320 IU/LB
vitamin E	Minimum	987 IU/LB
ORGANIC SELE	Minimum	10 PPM

### Ingredient List

Calcium carbonate, Salt, Magnesium Oxide, Potassium Chloride, Intellibond micro 5, Mineral oil, vitamin E supplement, Selenium Yeast, Tri Basic Copper, Manganese Oxide, Sodium Selenite, vitamin A Acetate, Calcium Iodate, Vitamin D3 supplement, Ethylene Diamine Dihydro Iodide.

**NET WT 50 LBS (22.67) KG**

SGHEIFER1

**Manufactured for: Trouw Nutrition**

**3422 Dutch Hollow Rd**

**Strykersville, NY, 14145**

10/23/2020



SGHEIFER1

CONSULT YOUR FEED NUTRITIONIST FOR FEEDING RATES.