



~ P C  
**Trouw Heifer Mineral**  
**2 Oz w Rumensin**



MEDICATED

Growing cattle fed in confinement (dairy and beef replacement heifers): For improved feed efficiency.

**ACTIVE DRUG INGREDIENT**

**Monensin (as Monensin sodium) . 1612.4 g/ton**

**Guaranteed Analysis:**

Calcium	(Min)	14	%
Calcium	(Max)	17	%
Salt	(Min)	22	%
Salt	(Max)	26	%
Magnesium	(Min)	9.3	%
Potassium	(Min)	3.5	%
Iodine	(Min)	46	PPM
Cobalt	(Min)	39	PPM
Copper	(Min)	1,244	PPM
Zinc	(Min)	3,315	PPM
Manganese	(Min)	2,488	PPM
Selenium	(Min)	34.8	PPM
Vitamin A	(Min)	156,643	IU/LB
Vitamin D	(Min)	73,526	IU/LB
Vitamin E	(Min)	1,968	IU/LB

**Ingredients: Calcium Carbonate, Salt, Magnesium Oxide, Calcium Sulfate, Potassium Chloride, Vitamin E Supplement, Mineral Oil, Selenium Yeast, Monocalcium Phosphate, Sodium Selenite, Manganous Oxide, Basic Copper Chloride, Zinc Hydroxychloride, Vitamin D Supplement, Vitamin A Supplement, Manganese Oxide, Manganese Protein, Soybean Oil, Cobalt Carbonate and Calcium Iodate.**

**CAUTION:** A withdrawal time has not been established for preruminating calves. Do not use in calves to be processed for veal. Do not allow horses or other equines access to formulation containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin medicated cattle and goat feed is safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of Monensin recommended in the feeding directions as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and amount of refusals fed should be taken into consideration to prevent Monensin overdosing. **LIMITATIONS:** Feed to mature reproducing beef cows and non-lactating dairy cattle. Feed as supplemental feed, either hand-fed in a minimum of 1 pound of feed or mixed in a total ration. For improved feed efficiency, feed continuously at a rate of 50 to 200 milligrams Monensin per head per day.

**See consultant for complete feeding instructions. Follow label directions:**

**The addition to feed of higher concentrations of this premix containing selenium is not permitted.**

**Manufactured For:**

**Trouw Nutrition**

**Strykersville, NY 14145**

**Net Weight 50 lbs (22.67 kg)**

**TN-HF2R**

**MEDICATED**